

# Primitive Neurectodermal Tumour of Kidney – An Uncommon Tumour At a Rare Site

**Sunil R<sup>a</sup>, S Vasudevan<sup>a</sup>, Laila Rani<sup>b</sup>, Praveen Gopi<sup>a</sup>, Anu Mary<sup>b</sup>**

a. Department of Urology, Government Medical College, Trivandrum;

b. Department of Pathology, Government Medical College, Trivandrum \*

## ABSTRACT

Published on 27<sup>th</sup> September 2017

A case of Primitive Neuroectodermal Tumour (PNET) of the kidney in a 31-year-old gentleman is presented. Few cases are reported in the literature with a variable, nonspecific presentation and an aggressive behaviour. In our case, a radical nephrectomy, lymphadenectomy and IVC repair was performed. The surgical specimens were formalin-fixed and paraffin embedded. The sections were stained with routine H&E. Immunohistochemistry was performed. The immunohistochemical evaluation revealed CD-99positive, Membrane positivity, NSE positivity, Synaptophysin-focal positivity and EMA-negativity. The clinical presentation and the macroscopic aspect, together with the histological pattern, the cytological characteristic and the cellular immunophenotype addressed the diagnosis towards primary PNET of kidney. Since sometimes it is difficult to discriminate between PNET and Ewing's tumour, we reviewed the difficulties in differential diagnosis. These tumours have a common precursor but the stage of differentiation in which it is blocked is probably different. This could also explain their different biological behaviour and prognosis.

**Keywords:** Renal PNET, Ewing's Sarcoma

\*See End Note for complete author details

## CASE REPORT

A 31 year old gentleman presented with abdominal lump of one month duration, haematuria on and off 2 months duration which is gross, intermittent, painless with passage of clots and associated abdominal pain. On general examination patient is hypertensive. On physical examination revealed an abdominal lump of size 20\*15cm which is hard and immobile. MR Urogram revealed a large mass in right kidney having highly heterogeneous signals most of the mass shows intermediate low T2 signals and the mass involved renal artery and vein and extends over the IVC and psoas major (figure 1). Metastatic workup done came as negative (figure 2,3,4,5). Patient underwent radical nephrectomy and IVC repair on 02-08-2017 and the specimen of size 25\*20cm, histopathology report came as primitive neurectodermal tumour.

## DISCUSSION

PNET of the kidney was first reported by Seemayer and colleagues in 1975<sup>1</sup> and is exceedingly rare. It usually affects young adults at a median age of 28 years

and has a male predominance of 3:1.<sup>2,3</sup> The common presenting symptoms are flank or abdominal pain, mass in the abdomen, and haematuria.<sup>4</sup> Patients are usually asymptomatic until the tumour reaches a large size; the maximum diameter of such tumours is often 10 cm.<sup>5,6</sup> Systemic symptoms such as weight loss and fever may also occur.<sup>7</sup> Most of the patients present at an advanced stage with distant metastasis, which is in concordance with the aggressive behaviour of this tumour.<sup>2,8</sup> Common sites of metastasis include the lung, liver, and bone.<sup>4</sup>

The imaging characteristics of renal PNET are often nonspecific and overlap with those of other renal tumours, such as renal cell carcinoma, Wilms tumor, neuroblastoma, lymphoma, and desmoplastic small round cell tumor.<sup>9</sup> Renal PNETs appear hypoechoic, isoechoic, and / or hyperechoic to the adjacent renal parenchyma on ultrasound and show increased vascularity on Doppler imaging. CT scan shows a large heterogeneous mass with areas of hemorrhage or necrosis.<sup>10</sup> On magnetic resonance imaging (MRI), the tumor appears as a lobulated isointense and / or hypointense mass on T1-weighted images and as a heterogeneous

*Cite this article as:* Sunil R, Vasudevan S, Rani L, Praveen P, Mary A. Primitive Neurectodermal Tumour of Kidney – An Uncommon Tumour At a Rare Site. Kerala Medical Journal. 2017 Sep 27;10(3):121–4.

### Corresponding Author:

Dr Vasudevan S, Professor, Department of Urology, Medical College, Trivandrum; Chief Editor, Kerala Medical Journal  
Email: periamana@gmail.com




Figure 1. MR Urogram - A large mass in right kidney having highly heterogeneous signals most of the mass shows intermediate low T2 signals and the mass involved renal artery and vein and extends over the IVC and psoas major

to hyperintense mass on T2-weighted images, with heterogeneous contrast enhancement.<sup>4,10</sup> MRI and CT also help to evaluate renal vein and inferior vena caval involvement. Technetium-99m scintigraphy is useful for the detection of bone metastases.<sup>4</sup>

Diagnosis of renal PNET is challenging. Although radiological features may be suggestive, biopsy with immunohistochemistry is required to confirm the diagnosis. Renal PNET is characterized by small uniform round cells with dark nuclei, ill-defined cytoplasmic borders, and poorly formed rosette-like structures.<sup>7</sup> The histopathologic features overlap with other small round blue cell tumors like neuroblastoma, desmoplastic small round cell tumor, and lymphoma. Immunohistochemistry and molecular studies play a crucial role

in differentiating these tumors. PNET shows strong positivity for MIC-2 gene product and CD-99 over the membrane of tumor cells, which is seen in more than 90% of renal PNET cases.<sup>5</sup> It is also positive for different neural biomarkers such as S-100, Leu 7, and NSE.<sup>11</sup> The Immunohistochemistry findings are further supported by the identification of a characteristic EWSR1/FLI1 fusion product that results from a t(11;22) (q24/q22;q12) translocation. This translocation is identified in 90% of cases and unequivocally confirms the diagnosis.<sup>5,12</sup> Among our case, IHC details were positive for CD99, NSE, Synaptophysin

Renal PNET appears to be a unique clinical entity that behaves more aggressively than PNET arising at other sites. As the tumor is highly aggressive, it is often




Figure 2. Gross specimen



Figure 3. Pathologic specimen




Figure 4. Microscopy

diagnosed in an advanced stage when it has already involved perinephric fat, hilar lymph nodes, renal veins, and the inferior vena cava. In more advanced stages, the tumor involves the liver, spleen, peritoneum, and lungs. The prognosis for renal ES/PNET is generally poor, with a 5-year disease-free survival of 45% to 55% in localized cases, whereas cases with an advanced stage at presentation have a median relapse-free survival of only 2 years.<sup>7,11,13</sup>

The treatment for renal ES/PNET is similar to that for ES/PNET elsewhere and includes surgery, chemotherapy, and radiation.<sup>4,8</sup> Surgical options include partial or total nephrectomy with cavotomy in cases of renal vein involvement.<sup>4,12</sup> The diagnosis of renal ES/PNET is often made postoperatively and hence chemotherapy is generally given as an adjuvant. The recommended chemotherapy regimen is vincristine, doxorubicin, and cyclophosphamide alternating with ifosfamide and etoposide given for a period of 1 year.<sup>14</sup> The role of radiotherapy is not clear, but it may be given in locally advanced disease and in those with positive margins. Despite aggressive therapy, the overall cure rate of renal PNET is only 20%.<sup>15</sup> Since the overall prognosis of this tumor is poor, an early and accurate diagnosis is crucial for the proper management of these aggressive tumors.

## END NOTE

### Author Information

1. Dr Sunil R, Senior Resident, Department of Urology, Government Medical College Trivandrum
2. Prof S Vasudevan, Professor, Department of Urology, Medical College, Trivandrum; Chief Editor, Kerala Medical Journal
3. Prof Laila Rani, Department of Pathology, Government Medical College Trivandrum



Figure 5. Technetium-99m scintigraphy bone scan normal study

4. Dr Praveen Gopi, Senior Resident, Department of Urology, Government Medical College, Trivandrum
5. Dr Anu Mary, Senior Resident, Department of Pathology, Government Medical College, Trivandrum

**Conflict of Interest:** None declared

**Editor's Remarks:** Primitive Neuroectodermal Tumors are very rare but aggressive tumors seen in younger patients often with florid manifestations. The clinical outcomes are often dismal.

## REFERENCES

1. Seemayer TA, Thelmo WL, Bolande RP, Wiglesworth FW. Peripheral neuroectodermal tumors. *Perspect Pediatr Pathol.* 1975;2(2):151–172.
2. Mohsin R, Hashmi A, Mubarak M, Sultan G, Shehzad A, Qayum A, Anwer Naqvi SA, Hassan Rizvi SA. Primitive neuroectodermal tumor/Ewing's sarcoma in adult uro-oncology: a case series from a developing country. *Urol Ann.* 2011;3(2):103–107.
3. Lee H, Cho JY, Kim SH, Jung DC, Kim JK, Choi HJ. Imaging findings of primitive neuroectodermal tumors of the kidney. *J Comput Assist Tomogr.* 2009;33(6):882–886.
4. Hari S, Jain TP, Thulkar S, Bakhshi S. Imaging features of peripheral primitive neuroectodermal tumours. *Br J Radiol.* 2008;81(972):975–983.
5. Wu YJ, Zhu YC, Chen H, Huang Y, Wei Q, Chen HJ, Xie X, Li X, Zhou Q, Yang YR, Zeng H. Primitive neuroectodermal tumor of the kidney with inferior vena cava tumor thrombus during pregnancy response to sorafenib. *Chin Med J (Engl)* 2010;123(15):2155–2158.
6. Ohgaki K, Horiuchi K, Mizutani S, Sato M, Kondo Y. Primary Ewing's sarcoma/primitive neuroectodermal tumor of the kidney that responded to low-dose chemotherapy with ifosfamide etoposide, and doxorubicin. *Int J Clin Oncol.* 2010;15(2):210–214.
7. Parham DM, Roloson GJ, Feely M, Green DM, Bridge JA, Beck-

with JB. Primary malignant neuroepithelial tumors of the kidney: a clinicopathologic analysis of 146 adult and pediatric cases from the National Wilms' Tumor Study Group Pathology Center. *Am J Surg Pathol.* 2001;25(2):133–146.

8. Ellinger J, Bastian PJ, Hauser S, Biermann K, Müller SC. Primitive neuroectodermal tumor: rare highly aggressive differential diagnosis in urologic malignancies. *Urology.* 2006;68(2):257–262.
9. Lalwani N, Prasad SR, Vikram R, Katabathina V, Shambhogue A, Restrepo C. Pediatric and adult primary sarcomas of the kidney: a cross-sectional imaging review. *Acta Radiol.* 2011;52(4):448–457.
10. Tappouni R, El-Kady R, Raman J, Then M, Sarwani N. Renal neuroectodermal tumour presenting with hematuria. *J Radiol Case Rep.* 2010;4(7):12–20.
11. Habermann H, Benesch M, Schips L, Pummera K, Ratschek M, Uggowitz MM, Urbanb C, Hubmera G. Findings and clinical course of a localized primitive peripheral neuroectodermal tumor of the kidney. *Urol Int.* 2003;71(3):319–321.
12. Fergany AF, Dhar N, Budd GT, Skacel M, Garcia JA. Primary extraosseous Ewing sarcoma of the kidney with level III inferior vena cava thrombus. *Clin Genitourin Cancer.* 2009;7(3):e95–e97.
13. Casella R, Moch H, Rochiltz C, Meier V, Seifert B, Mihatsch MJ, Gasser TC. Metastatic primitive neuroectodermal tumor of the kidney in adults. *Eur Urol.* 2001;39(5):613–617.
14. Grier HE, Kralo MD, Tarbell NJ, Link MP, Fryer CJ, Pritchard DJ, Gebhardt MC, Dickman PS, Perlman EJ, Meyers PA, Donaldson SS, Moore S, Rausen AR, Vietti TJ, Miser JS. Addition of ifosfamide and etoposide to standard chemotherapy for Ewing's sarcoma and primitive neuroectodermal tumor of bone. *N Engl J Med.* 2003;348(8):694–701.
15. Angel JR, Alfred A, Sahuja A, Sells RE, Zechlinski JJ. Ewing's sarcoma of the kidney. *Int J Clin Oncol.* 2010;15(3):314–318.